

Analysis on Symmetric and Locally Symmetric Spaces

(Multiplicities, Cohomology and Zeta functions)

Ulrich Bunke and Robert Waldmüller

1 Multiplicities of principal series

Let us first fix some notation. Let G be a semisimple Liegroup with finite center and $G \cong KAN$ be the Iwasawa decomposition of G , where K is a maximal compact subgroup, A a maximal $\mathbb{R}$ -split torus of G , and N is a nilpotent subgroup. Correspondingly, we have a decomposition of the Lie algebra: $\mathfrak{g} \cong \mathfrak{k} \oplus \mathfrak{a} \oplus \mathfrak{n}$. We denote by $P := MAN$ a minimal parabolic subgroup, where $M = Z_K(A)$ is the centraliser of K in A . For $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$, we define the character $a \mapsto a^\lambda := \exp(\lambda(\log a))$. Furthermore, let $\rho \in \mathfrak{a}^*$ denote $H \mapsto \frac{1}{2}\text{Tr}(ad(H)|_{\mathfrak{n}})$. For a finite dimensional representation (σ, V_σ) of M , we define $\sigma_\lambda : P \rightarrow GL(V_\sigma)$ by $man \mapsto a^{\rho-\lambda}\sigma(m)$.

Definition 1 *The induced representation $\pi^{\sigma_\lambda} := \text{Ind}_P^G(\sigma_\lambda)$ is called the principal series representation with parameter (σ, λ) .*

π^{σ_λ} acts on the space of sections of regularity ? of the G -equivariant bundle $V(\sigma, \lambda) := G \times_P V_{\sigma_\lambda}$. The space of sections will be denoted by $V_{\pi^{\sigma_\lambda}}^? := C^?(G/P, V(\sigma, \lambda))$, where ? belongs to $K - fin, \omega, \infty, L^2, -\infty$ or $-\omega$. Here, $K - fin$ means K -finite, ω real-analytic, ∞ smooth and L^2 square integrables. To be honest, $C^{K-fin}(G/P, V(\sigma, \lambda))$ is only a $(\mathfrak{g}, K)$ -module. Denote the $\mathbb{C}$ -linear dual of σ by σ^* . To define the hyperfunction and distribution valued sections, observe that $V(1, -\rho) \rightarrow G/P$ is the bundle of densities, and we have a G -equivariant pairing $V(\sigma, \lambda) \otimes V(\sigma^*, -\lambda) \rightarrow V(1, -\rho)$. Combining this pairing with integration, we have a pairing of spaces of sections.

Definition 2

$$\begin{aligned} C^{-\omega}(G/P, V(\sigma, \lambda)) &:= C^\omega(G/P, V(\sigma^*, -\lambda))^* \\ C^{-\infty}(G/P, V(\sigma, \lambda)) &:= C^\infty(G/P, V(\sigma^*, -\lambda))^*. \end{aligned}$$


$C^{L^2}(G/P, V(\sigma, \lambda))$ is a G -Banach representation. If $\Re(\lambda) = 0$, it is unitary in a natural way.

Example We consider $G = SL(2, \mathbb{R})$. The Iwasawa components are

$$K = \left\{ \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \mid \phi \in [0, 2\pi) \right\} \quad A = \left\{ \begin{pmatrix} t^{\frac{1}{2}} & 0 \\ 0 & t^{-\frac{1}{2}} \end{pmatrix} \mid t \in (0, \infty) \right\}$$

$$N = \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mid x \in \mathbb{R} \right\}$$

In this case, we have $\mathfrak{a} \cong \mathbb{R}H$, where $H = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix}$. Calculating $\rho(H) = \frac{1}{2}$, we obtain that $\rho \cong \frac{1}{2}$. Since $M \cong \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \right\} \cong \mathbb{Z}/2\mathbb{Z}$, $\hat{M} = \{1, \theta\}$ consists of two elements. Furthermore, $G/P = K/M \cong S^1$. Choosing $\sigma = 1$, the principal series representations are parametrized by $\lambda \in \mathbb{C}$. For generic λ , the representation is irreducible.



D_{2n+1}^+ is called the holomorphic/antiholomorphic discrete series, and F_{2n+1} is the $2n+1$ -dimensional irreducible representation of $SL(2, \mathbb{R})$. The unitary principal series representations, the complementary series representations, $D_{2n+1}^\pm$ and F_1 are unitary. Note that $\pi^{1 \pm \frac{1}{2}}$ is a non-trivial extension and therefore not unitary.

Let $\Gamma \subset G$ be a cocompact, torsion-free discrete subgroup, $X := G/K$, $Y := \Gamma \backslash X$ and ${}^\Gamma V_{\pi^{\sigma\lambda}}^{-\omega} := \{\phi \in V_{\pi^{\sigma\lambda}}^{-\omega} | \pi^{\sigma\lambda}(\gamma)\phi = \phi \ \forall \gamma \in \Gamma\}$, the space of Γ -invariant hyperfunction vectors.

Lemma 1 (Frobenius reciprocity)

$$\begin{aligned} {}^\Gamma V_{\pi^{\sigma\lambda}}^{-\omega} &\cong \text{Hom}_G(V_{\pi^{\sigma^*-\lambda}}^{-\omega}, C^\infty(\Gamma \backslash G)) \\ \phi &\mapsto (f \mapsto (g \mapsto \phi(\pi^{\sigma^*-\lambda}(g)f))) \\ (f \mapsto (\psi(f)(\Gamma 1))) &\leftarrow \psi \end{aligned}$$

Since Γ is cocompact, we have a decomposition

$$C^\infty(\Gamma \backslash G) \cong \widehat{\bigoplus_{(\pi, V_\pi) \in \hat{G}_u} m_\Gamma(\pi) V_\pi^\infty},$$

where $\hat{G}_u$ is the unitary dual of G , i.e. the set of equivalence classes of irreducible unitary representations of G . As a consequence, if $\pi^{\sigma\lambda}$ is unitary and $\lambda \neq 0$, then it is irreducible and therefore $m_\Gamma(\pi^{\sigma\lambda}) = \dim({}^\Gamma V_{\pi^{\sigma\lambda}}^{-\omega})$. Moreover, if $\pi^{\sigma\lambda}$ is irreducible and not unitary, then ${}^\Gamma V_{\pi^{\sigma\lambda}}^{-\omega} = \{0\}$.

We now return to our example and connect ${}^\Gamma V_{\pi^{\sigma\lambda}}^{-\omega}$ with the eigenvalues of the Laplacian on Y . As K -homogenous bundles, $V(\sigma, \lambda) = G \times_P V_{\sigma\lambda} = K \times_M V_\sigma$, and for $\sigma = 1$, $K \times_M V_1 \cong S^1 \times \mathbb{C}$. Denote by $f \in C^\infty(K \times_M V_1)$ the unique K -invariant section with $f(1) = 1$. For any choice of λ , this corresponds to a real-analytic section f_λ of $V(1, \lambda)$.

Using the reciprocity homomorphism $i : V_{\pi^{1\lambda}}^{-\omega} \rightarrow \text{Hom}_G(V_{\pi^{1-\lambda}}^\omega, C^\infty(G))$ for trivial Γ , we define the Poisson transformation

$$i.(f_{-\lambda}) : V_{\pi^{1\lambda}}^{-\omega} \rightarrow C^\infty(G).$$

Now, any $\phi \in V_{\pi^{1\lambda}}^{-\omega}$ defines $i_\phi(f_{-\lambda}) \in C^\infty(X)$. The Casimir-operator Ω of $\mathfrak{g}$ acts on $\pi^{1\lambda}$ by a scalar, $\pi^{1\lambda}(\Omega) = \mu(\lambda) := \frac{1}{4} - \lambda^2$, while $\Omega|_{C^\infty(X)} = \Delta_X$, the Laplace operator. Therefore, $i_\phi(f_{-\lambda}) \in \text{Ker}(\Delta_X - \mu(\lambda))$.

Theorem 1 (Helgasson [9]) *For $\lambda \notin \{-\frac{1}{2}, -\frac{3}{2}, -\frac{5}{2}, \dots\}$, the Poisson transformation*

$$i.(f_{-\lambda}) : V_{\pi^{1\lambda}}^{-\omega} \rightarrow \text{Ker}(\Delta_X - \mu(\lambda))$$

is an isomorphism.

To calculate $H^*(\Gamma, V_{\pi^{1\lambda}}^{-\omega})$, we need the following classical theorems.

Theorem 2 *Let M be a C^ω -manifold, $E, F \rightarrow M$ C^ω -vector bundles and $A : C^\infty(M, E) \rightarrow C^\infty(M, F)$ an elliptic operator with C^ω -coefficients. If M is non-compact, then A is surjective.*

Theorem 3 ([1], Lemma 2.4, 2.6) *Let a discrete group U act properly on a space M and let $\mathcal{F}$ be a soft or flabby U -equivariant sheaf on M . Then $H^i(U, \mathcal{F}(M)) = 0$ for all $i \geq 1$.*

Combining Theorem 1 and Theorem 2, we get a short exact sequence for $\lambda \notin \{-\frac{1}{2}, -\frac{3}{2}, -\frac{5}{2}, \dots\}$

$$0 \rightarrow V_{\pi^1 \lambda}^{-\omega} \xrightarrow{i \cdot (f_{-\lambda})} C^\infty(X) \xrightarrow{\Delta_X - \mu(\lambda)} C^\infty(X) \rightarrow 0.$$

By Theorem 3, this is a Γ -acyclic resolution of $V_{\pi^1 \lambda}^{-\omega}$, so for $\lambda \notin -\frac{1}{2} - \mathbb{N}_0$, we get

i	$H^i(\Gamma, V_{\pi^1 \lambda}^{-\omega})$
0	$\ker(\Delta_Y - \mu(\lambda))$
1	$\operatorname{coker}(\Delta_Y - \mu(\lambda))$

Since $(\Delta_Y - \mu(\lambda))$ is an elliptic operator, both spaces are finite dimensional. To compute ${}^\Gamma V_{\pi^1 \lambda}^{-\omega}$ for $\lambda \in -\frac{1}{2} - \mathbb{N}_0$, we need $H^i(\Gamma, V_{\pi^1 \lambda}^{-\omega})$ for $i \geq 1, \lambda \in \frac{1}{2} + \mathbb{N}$ as well.

In the case of $\lambda \in \frac{1}{2} + \mathbb{N}$, we have the exact sequence

$$0 \rightarrow F_{2n+1} \rightarrow V_{\pi^{1_{n+\frac{1}{2}}}}^{-\omega} \rightarrow D_{2n+1}^{-\omega} \rightarrow 0$$

Since in this case $\mu(\lambda) = \frac{1}{4} - \lambda^2$ is negative, $\Delta_Y - \mu(\lambda)$ is a positive operator on a compact manifold and therefore injective, so ${}^\Gamma V_{\pi^1 \lambda}^{-\omega}$ is trivial. Using the long exact sequence

$$\dots \rightarrow H^i(\Gamma, F_{2n+1}) \rightarrow H^i(\Gamma, V_{\pi^{1_{n+\frac{1}{2}}}}^{-\omega}) \rightarrow H^i(\Gamma, D_{2n+1}^{-\omega}) \rightarrow \dots$$

and the topological result

i	$\dim(H^i(\Gamma, F_{2n+1}))$	$\dim(H^i(\Gamma, F_1))$
0	0	1
1	$(2g-2)(2n+1)$	2g
2	0	1

where g denotes the genus of Y , we get

i	$\dim(H^i(\Gamma, F_{2n+1}))$	$\dim(H^i(\Gamma, V_{\pi^1 \lambda}^{-\omega}))$	$\dim(H^i(\Gamma, D_{2n+1}^{-\omega}))$
0	0	0	$(2g-2)(2n+1)$
1	$(2g-2)(2n+1)$	0	0
2	0	0	0

For $\lambda = 1/2$, we have that $\dim(\ker(\Delta_Y)) = \dim(\operatorname{coker}(\Delta_Y)) = 1$, so we get

i	$\dim(H^i(\Gamma, F_1))$	$\dim(H^i(\Gamma, V_{\pi^{1/2}}^{-\omega}))$	$\dim(H^i(\Gamma, D_1^{-\omega}))$
0	1	1	2g
1	2g	1	2
2	1	0	0

Here, the long exact sequence gave us that $\dim(H^1(\Gamma, D_1^{-\omega}))$ is either 1 or 2 and since $D_1^{-\omega}$ is the sum of two conjugate isomorphic G submodules, it has to be even. For $\lambda \in -1/2 - \mathbb{N}_0$, we have a long exact sequence

$$\dots \rightarrow H^i(\Gamma, D_{2n+1}^{-\omega}) \rightarrow H^i(\Gamma, V_{\pi^{-1}\lambda}^{-\omega}) \rightarrow H^i(\Gamma, F_{2n+1}) \rightarrow \dots$$

Using the result for positive λ , we obtain immediately for $\lambda \in -\frac{1}{2} - \mathbb{N}$

i	$\dim(H^i(\Gamma, D_{2n+1}^{-\omega}))$	$\dim(H^i(\Gamma, V_{\pi^{-1}\lambda}^{-\omega}))$	$\dim(H^i(\Gamma, F_{2n+1}))$
0	$(2n+1)(2g-2)$	$(2n+1)(2g-2)$	0
1	0	$(2n+1)(2g-2)$	$(2n+1)(2g-2)$
2	0	0	0

In the case of $\lambda = -1/2$, it is

i	$\dim(H^i(\Gamma, D_1^{-\omega}))$	$\dim(H^i(\Gamma, V_{\pi^{-1}\frac{1}{2}}^{-\omega}))$	$\dim(H^i(\Gamma, F_1))$
0	$2g$	$2g$	1
1	2	$2g+1$	$2g$
2	0	1	1

Everything except $\dim(H^0(\Gamma, V_{\pi^{-1}\frac{1}{2}}^{-\omega}))$ is determined by the long exact sequence. For $\dim(H^0(\Gamma, V_{\pi^{-1}\frac{1}{2}}^{-\omega}))$, we could have $2g$ or $2g+1$, but all invariants are in the submodule $D_1^{-\omega}$ because otherwise we would get an embedding of $V_{\pi^{-1}\frac{1}{2}}^{\omega}$ into $L^2(\Gamma \backslash G)$ which is impossible, since $V_{\pi^{-1}\frac{1}{2}}^{\omega}$ is not unitary.

In all these cases, the cohomology groups are finite dimensional and $\mathcal{X}(\Gamma, V_{\pi^{-1}\lambda}^{-\omega}) = 0$.

Let us now consider the general case, so let $G_{\mathbb{C}}$ be a connected reductive group over $\mathbb{C}$, $G_{\mathbb{R}}$ a real form, $K \subset G_{\mathbb{R}}$ maximal compact, and $\Gamma \subset G_{\mathbb{R}}$ be cocompact, torsion-free and discrete. Denote by $Mod(\mathfrak{g}, K)$ the category of $(\mathfrak{g}, K)$ modules and by $\mathcal{HC}(\mathfrak{g}, K) \subset Mod(\mathfrak{g}, K)$ the subcategory of Harish-Chandra modules. Recall that a module is called Harish-Chandra if it is finitely generated over the universal enveloping algebra $\mathcal{U}(\mathfrak{g})$ and admissible, i.e. $\forall \gamma \in \hat{K}$, $\dim(V(\gamma)) < \infty$, where $V(\gamma)$ is the γ -isotypical component of V . For a G module V denote the K -finite submodule by $V_{K-fin} = \{v \in V \mid \dim(\text{span}(Kv)) < \infty\}$. For $V \in \mathcal{HC}(\mathfrak{g}, K)$ denote by $\tilde{V} := Hom(V, \mathbb{C})_{K-fin}$ the dual in $\mathcal{HC}(\mathfrak{g}, K)$.

Definition 3 For a Harish-Chandra module V , the maximal globalisation of V is $MG(V) := Hom_{(\mathfrak{g}, K)}(\tilde{V}, C^{\infty}(G)_{K-fin})$.

Thus, the maximal globalisation $MG(V)$ is a particular continuous representation of G whose K -finite part is isomorphic to V .

For a Harish-Chandra module V , define the hyperfunction vectors of V to be $V^{-\omega} := (\tilde{V}_B^{\omega})^*$, the topological dual of the analytic vectors in any Banach globalisation of the dual representation.

Theorem 4 (Schmid [15]) $MG(V) \cong V^{-\omega}$.

Theorem 5 (Bunke/Olbrich [2], 1.4) For $V \in \mathcal{HC}(\mathfrak{g}, K)$,

$$H^*(\Gamma, MG(V)) \cong \bigoplus_{(\pi, V_\pi) \in \hat{G}_u} m_\Gamma(\pi) Ext_{(\mathfrak{g}, K)}^*(\tilde{V}, V_{\pi, K-fin}).$$

For the proof of this theorem, we need the generalisation of Theorem 3

$$H^i(\Gamma, C^\infty(G)_{K-fin}) = 0 \quad \forall i \geq 1$$

and the following generalisation of Theorem 1.

Theorem 6 (Kashiwara-Schmidt [12])

$$Ext_{(\mathfrak{g}, K)}^i(\tilde{V}, C^\infty(G)_{K-fin}) = \begin{cases} MG(V), & i = 0 \\ 0, & i > 0 \end{cases}$$

Now, let us chose a projective resolution $P. \rightarrow \tilde{V} \rightarrow 0$ of $\tilde{V}$ in $Mod(\mathfrak{g}, K)$. By Theorem 6, $Hom_{(\mathfrak{g}, K)}(P., C^\infty(G)_{K-fin})$ resolves $MG(V)$ Γ -acyclically, therefore $Hom_{\mathfrak{g}, K}(P., C^\infty(\Gamma \backslash G)_{K-fin})$ calculates $H^*(\Gamma, MG(V))$. Finally, an analytic argument ([2], Lemma 3.1) shows that only a finite part of the decomposition

$$C^\infty(\Gamma \backslash G)_{K-fin} = \widehat{\bigoplus_{(\pi, V_\pi) \in \hat{G}_u} m_\Gamma(\pi) V_{\pi, K-fin}^\infty}$$

contributes to $Hom_{(\mathfrak{g}, K)}(P., C^\infty(\Gamma \backslash G)_{K-fin})$.

2 The Selberg Zeta function

In this section we will additionally assume that the real rank of G is equal to one. A group element $g \in G$ is called hyperbolic if there exist $m_g \in M$, $a_g \in A$ such that g is conjugate in G to $m_g a_g$ with $a_g^\rho > 1$. If g is hyperbolic, a_g is unique, and m_g is unique up to conjugation in M . Note that if $\gamma \in \Gamma$, $\gamma \neq 1$, then γ is hyperbolic.

For a hyperbolic $g \in G$, we define

$$Z(g, \sigma, \lambda) := \prod_{k=0}^{\infty} det(1 - \sigma_{\lambda-2\rho}(m_g a_g) \otimes S^k(Ad(m_g a_g)|_{\overline{\mathfrak{n}}})),$$

where $\sigma_{\lambda-2\rho}$ denotes the representation of P as in section 1 and S^k is the k 'th symmetric power.

In the example of $G = SL(2, \mathbb{R})$, a hyperbolic g is conjugate to $m_g \begin{pmatrix} t^{1/2} & 0 \\ 0 & t^{-1/2} \end{pmatrix}$, $t > 1$. Therefore, we get

$$Z(g, 1, \lambda) = \prod_{k=0}^{\infty} (1 - t^{-\lambda-k-1/2}).$$

Definition 4 The conjugacy class $[\gamma] \in C\Gamma$ is called *primitive*, if $[\gamma]$ is not of the form $[\gamma] = [\gamma'^n]$ for some $\gamma' \in \Gamma, n > 1$.

Definition 5 The Selberg Zeta function is defined for $\Re(\lambda) > \rho$ by the converging product

$$Z(\Gamma, \sigma, \lambda) = \prod_{\substack{[\gamma] \in C\Gamma \\ [\gamma] \text{ primitive}}} Z(\gamma, \sigma, \lambda).$$

Theorem 7 (Fried [8]) $Z(\Gamma, \sigma, \lambda)$ has a meromorphic continuation to $\mathfrak{a}_{\mathbb{C}}^*$.

The proof uses Ruelles thermodynamic formalism. The disadvantage of this method of proof is that it doesn't give any information about the singularities of the continuation or about a functional equation.

We use the Selberg trace formula to calculate the singularities of the Zeta function and to determine the functional equation. To this end, we introduce the logarithmic derivative

$$L(\Gamma, \sigma, \lambda) := \frac{Z'(\Gamma, \sigma, \lambda)}{Z(\Gamma, \sigma, \lambda)},$$

considered as a 1-form on $\mathfrak{a}_{\mathbb{C}}^*$. Denote by X^d the dual of the symmetric space X , i.e.

X	H^m	$\mathbb{C}H^m$	$\mathbb{H}H^m$	CaH^2
X^d	S^n	$\mathbb{C}P^m$	$\mathbb{H}P^m$	CaP^2

In the case of $n := \dim X$ odd, we assume that either σ is irreducible and the isomorphism class of σ is fixed under the action of the Weyl group, or $\sigma = \sigma' \oplus \sigma'^w$ with w the non-trivial element of the Weyl group $W(\mathfrak{g}, \mathfrak{a}) \cong \mathbb{Z}/2\mathbb{Z}$ and σ' irreducible.

The positive root system of $(\mathfrak{g}, \mathfrak{a})$ is either of the form $\{\alpha\}$ or $\{\frac{\alpha}{2}, \alpha\}$. We call α the long root and identify $\mathfrak{a}_{\mathbb{C}}^* \cong \mathbb{C}$ sending α to 1. Define the root vector $H_{\alpha} \in \mathfrak{a}$ corresponding to a positive root α by the condition that

$$\lambda(H_{\alpha}) = \frac{\langle \lambda, \alpha \rangle}{\langle \alpha, \alpha \rangle} \quad \forall \lambda \in \mathfrak{a}^*,$$

where $\langle \cdot, \cdot \rangle$ denotes an $Ad(G)$ -invariant inner product on $\mathfrak{g}$. For $\sigma \in \hat{M}$, we define $\epsilon_{\alpha}(\sigma) \in \{0, \frac{1}{2}\}$ by the condition $e^{2\pi i \epsilon_{\alpha}(\sigma)} = \sigma(\exp(2\pi i H_{\alpha}))$ and $\epsilon_{\sigma} \in \{0, \frac{1}{2}\}$ by requiring $\epsilon_{\sigma} \equiv |\rho| + \epsilon_{\alpha}(\sigma) \pmod{\mathbb{Z}}$.

Theorem 8 (Bunke/Olbrich [6]) *There exists a virtual elliptic operator $A_X(\sigma)$ and a corresponding operator $A_{X^d}(\sigma)$, such that in the case of n even*

$$\frac{L(\Gamma, \sigma, \lambda)}{2\lambda} = {}'Tr'(\lambda^2 + A_Y(\sigma))^{-1} \quad (1)$$

$$- \frac{(-1)^{\frac{n}{2}} \text{vol}(Y) \pi}{\text{vol}(X^d)} \frac{P_\sigma(\lambda)}{2\lambda} \begin{cases} \tan(\pi\lambda) & \text{if } \epsilon_\sigma = \frac{1}{2} \\ -\cot(\pi\lambda) & \text{if } \epsilon_\sigma = 0 \end{cases} \quad (2)$$

$$- \frac{(-1)^{\frac{n}{2}} \text{vol}(Y)}{\text{vol}(X^d)} {}'Tr'(\lambda^2 - A_{X^d}(\sigma))^{-1} \quad (3)$$

and in the case of n odd,

$$\begin{aligned} \frac{L(\Gamma, \sigma, \lambda)}{2\lambda} &= {}'Tr'(\lambda^2 + A_Y(\sigma))^{-1} \\ &+ (-1)^{\frac{n+1}{2}} \frac{\text{vol}(Y) \pi}{\text{vol}(X^d)} \frac{P_\sigma(\lambda)}{2\lambda} \end{aligned}$$

In the theorem, P_σ is some polynomial depending on σ . Since $(\lambda^2 + A_Y^2(\sigma))^{-1}$ is not trace-class, we have to use a regularised trace $'Tr'$. We refer to [6], 3.2. for a discussion of the regularisation.

From this formula, it follows that $L(\Gamma, \sigma, \lambda)$ is meromorphic.

In the case of n even, (2) is odd and (3) is even. The poles of the two terms cancel for $\lambda > 0$, add up for $\lambda < 0$ and $I := 2\lambda((2) + (3))$ is regular at zero ([6], 3.2.3). Hence, the set of poles of I is $\epsilon_\sigma - \mathbb{N}$.

For A an operator on some Hilbert space, $'Tr'(\lambda - A)^{-1}$ has first order poles at the eigenvalues of A with $\text{res } (2\lambda {}'Tr'(\lambda^2 - A)^{-1})$ equal to the dimension of the corresponding eigenspace, hence


$$\text{res}_{-\epsilon_\sigma - k} I = - \frac{(-1)^{\frac{n}{2}} 2\text{vol}(Y)}{\text{vol}(X^d)} \dim E_{A_{X^d}(\sigma)}(-\epsilon_\sigma - k)^2.$$

It follows from Hirzebruch proportionality that $\frac{\text{vol}(Y)}{\text{vol}(X^d)} \in \mathbb{Z}$ ([6], Prop. 3.14), so the residues of I are integral.

(1) has first order poles at $\pm i s$ with residue $\dim E_{A_Y(\sigma)}(s^2)$, where s^2 is a non-zero eigenvalue of $A_Y(\sigma)$ and if zero is an eigenvalue of $A_Y(\sigma)$, there is an additional pole at zero with residue $2\dim \ker A_Y(\sigma)$.

In the case of n odd, the only poles are those coming from the first term.


Note that all the residues of L are integral.



We'll now come back to the example of $G = SL(2, \mathbb{R})$, $\sigma = 1$. The above theorem specializes to

$$\frac{L(\Gamma, \sigma, \lambda)}{2\lambda} := {}'Tr'(\Delta_Y - \frac{1}{4} + \lambda^2)^{-1} + \frac{2g-2}{2} {}'Tr'(\Delta_{S^2} + \frac{1}{4} - \lambda^2)^{-1} + \frac{2g-2}{2} \frac{\pi \tan(\lambda\pi)}{\lambda},$$

where g is the genus of $\Gamma \backslash X$. The eigenvalues of Δ_{S^2} are $\frac{n^2}{4}$, $n \in \mathbb{N}$ with $mult(n^2/4) = 2n+1$. Using this, we get the following picture for $ord(Z(\Gamma, 1, \lambda))$.



Theorem 9 (Bunke/Olbrich, conjectured by Patterson [1])

For $\lambda \neq 0$ ¹,

$$\text{ord}_\lambda Z(\Gamma, \sigma, \lambda) = -\mathcal{X}'(\Gamma, V_{\pi\sigma_\lambda}^{-\omega})$$

where $\mathcal{X}'(\Gamma, V_{\pi\sigma_\lambda}^{-\omega}) = \sum_{p=0}^{\infty} (-1)^p p \dim H^p(\Gamma, V_{\pi\sigma_\lambda}^{-\omega})$.

Proof: In the example, the proof is just by inspection. In the general case, use Theorem 5 to obtain

$$\begin{aligned} & H^p(\Gamma, V_{\pi\sigma_\lambda}^{-\omega}) \\ &= \bigoplus_{(\pi, V_\pi) \in \hat{G}_u} m_\Gamma(\pi) \text{Ext}_{(\mathfrak{g}, K)}^p(V_{\pi\sigma_\lambda^*, K-\text{fin}}, V_{\pi, K-\text{fin}}) \\ &= \bigoplus_{(\pi, V_\pi) \in \hat{G}_u} m_\Gamma(\pi) \text{Ext}_{(\mathfrak{g}, K)}^p(V_{\pi^*, K-\text{fin}}, V_{\pi\sigma_\lambda, K-\text{fin}}) \\ &= \bigoplus_{(\pi, V_\pi) \in \hat{G}_u} m_\Gamma(\pi) \text{Hom}_{MA}(H_p(\mathfrak{n}, V_{\pi^*}) \oplus H_{p-1}(\mathfrak{n}, V_{\pi^*}), V_{\sigma_\lambda}) \\ &= \bigoplus_{(\pi, V_\pi) \in \hat{G}_u} m_\Gamma(\pi) \text{Hom}_{MA}([H^{n-1-p}(\mathfrak{n}, V_{\pi^*, K-\text{fin}}) \\ &\quad \oplus H^{n-p}(\mathfrak{n}, V_{\pi^*, K-\text{fin}})] \otimes \Lambda^{n-1} \mathfrak{n}, V_{\sigma_\lambda}) \\ &= \bigoplus_{(\pi, V_\pi) \in \hat{G}_u} m_\Gamma(\pi) \left[((H^p(\mathfrak{n}, V_{\pi, K-\text{fin}}) \oplus H^{p-1}(\mathfrak{n}, V_{\pi, K-\text{fin}})) \otimes V_{\sigma_\lambda^*})^{MA} \right]^* \end{aligned}$$

Setting

$$\mathcal{X}(\pi, \sigma, \lambda) := \sum_{p=0}^{\infty} (-1)^p \dim(H^p(\mathfrak{n}, V_{\pi, K-\text{fin}}) \otimes V_{\sigma_\lambda^*})^{MA},$$

the following theorem finishes the proof.

Theorem 10 (Juhl, [10], [11])

$$\text{ord}_\mu Z(\Gamma, \sigma, \lambda) = (-1)^{n-1} \sum_{(\pi, V_\pi) \in \hat{G}_u} m_\Gamma(\pi) \mathcal{X}(\pi, \sigma, \mu).$$

3 Generalisations

There are two obvious directions in which the theory could be generalised. First, one could try to generalise the group G . Unfortunately, we do not know of a satisfactory general definition of a Zeta function for groups of higher rank, but see [13], [14], [7] for special cases.

The other direction is to weaken the conditions on Γ . In [3] and [4], the theory is generalised to the case of subgroups $\Gamma \subset G$ of finite covolume. Examples of infinite covolume (convex cocompact) are considered in [5]

¹For $\lambda = 0$ see [1]

References

- [1] U. Bunke and M. Olbrich, *Gamma-cohomology and the Selberg zeta function* J. reine angew. Math. 467 (1995)
- [2] U. Bunke and M. Olbrich, *Cohomological Properties of the Canonical Globalizations of Harish-Chandra Modules* Annals of Global Analysis and Geometry 15 (1997)
- [3] U. Bunke and M. Olbrich, *Resolutions of distribution globalizations of Harish-Chandra modules and cohomology*, J. reine angew. Math. 497 (1998)
- [4] U. Bunke and M. Olbrich, *Fuchsian groups of the second kind and representations carried by the limit set*, Invent. math. 127 (1997)
- [5] U. Bunke and M. Olbrich, *Group cohomology and the singularities of the Selberg Zeta function associated to a Kleinian group*, Annals of mathematics 149 (1999)
- [6] U. Bunke and M. Olbrich, *Selberg Zeta and Theta Functions*, Math. Res. 83, Akademie Verlag, Berlin 1995
- [7] A. Deitmar, *Geometric zeta functions of locally symmetric spaces*, Amer. J. Math. 122 (2000)
- [8] D. Fried, *The zeta functions of Ruelle and Selberg I*, Ann. Sci. Ec. Norm. Sup. (4) 19 (1986)
- [9] S. Helgason, *Groups and Geometric Analysis*, Academic Press, 1984
- [10] A. Juhl, *Zeta-Funktionen, Index-Theorie und hyperbolische Dynamik*, Habilitationsschrift, Humboldt-Universität zu Berlin, 1995
- [11] A. Juhl, *Cohomological Theory of Dynamical Zeta Functions*, Birkhäuser, Basel 2001
- [12] M. Kashiwara and W. Schmid *Quasi-equivariant $\mathcal{D}$ -modules, equivariant derived category and representations of reductive Lie groups*, in J.L. Brylinski, R. Brylinski, V. Guillemin and V. Kac (eds.), *Lie Theory and Geometry. In Honor of Bertram Kostant*, Progr. in Math. 123, 1994
- [13] H. Moscovici and R. Stanton, *Eta invariants of Dirac operators on locally symmetric manifolds*, Invent. Math. 95 (1989)
- [14] H. Moscovici and R. Stanton, *R-torsion and zeta functions for locally symmetric manifolds*, Invent. Math. 105 (1991)
- [15] W. Schmid, *Boundary value problems for group invariant differential equations*, Soc. Math. de France, Asterisque, hors serie(1985)